

Co-occurring Mental Illness & Substance-Related Disorders

BIBLIOGRAPHY

- Ahmadi, J., Toobaee, S., Kharras, M., & Radmehr, M. (2003). Psychiatric disorders in opioid dependants. *International Journal of Social Psychiatry*, 49(3), 185–191.
- Albanese, M. J., & Pies, R. (2004). The bipolar patient with comorbid substance use disorder: Recognition and management. *CNS Drugs*, 18(9), 585–596.
- Alford DP, LaBelle CT, Richardson JM, O’Connell JJ, Hohl CA, Cheng DM, Samet JH. (2007). Treating homeless opioid dependent patients with buprenorphine in an office-based setting. *Journal of General Internal Medicine*, 22(2), 171–176.
- Bachman, S. S., Drainoni, M.-L., & Tobias, C. (2004). Medicaid managed care, substance abuse treatment, and people with disabilities: Review of the literature. *Health & Social Work*, 29(3), 189–196.
- Blow, F. C., Barry, K. L., BootsMiller, B. J., Copeland, L. A., McCormick, R., & Visnic, S. (1998). Longitudinal assessment of inpatient use and functioning of seriously mentally ill veterans with and without co-occurring substance use disorders. *Journal of Psychiatric Research*, 32(5), 311–319.
- Bogenschutz MP, Scott Tonigan J, Pettinati HM. (2009). Effects of alcoholism typology on response to naltrexone in the COMBINE study. *Alcohol Clinical Experimental Research*, 33(1), 10–18.
- Bond GR, Drake RE, Becker DR. (2008). An update on randomized controlled trials of evidence-based supported employment. *Psychiatric Rehabilitation Journal*, 31(4), 280–290.
- Boss EF, Zevin B. (2008), Neurobiology of Addiction. Presentation, National HCH Conference and Policy Symposium, Phoenix, AZ. <http://www.nhchc.org/2008conference/workshops/77.html>
- Bradizza, C. M., Stasiewicz, P. R., & Paas, N. D. (2006). Relapse to alcohol and drug use among individuals diagnosed with co-occurring mental health and substance use disorders: A review. *Clinical Psychological Review*, 26(2), 162–178.
- Brady, K. T., & Sinha, R. (2005). Co-occurring mental and substance use disorders: the neurobiological effects of chronic stress. *American Journal of Psychiatry*, 162(8), 1483–1493.
- Brady, K. T., & Verduin, M. L. (2005). Pharmacotherapy of comorbid mood, anxiety, and substance use disorders. *Substance Use & Misuse*, 40(13–14), 2021–2041, 2043–2048.
- Burns, L., Teesson, M., & O’Neill, K. (2005). The impact of comorbid anxiety and depression on alcohol treatment outcomes. *Addiction*, 100(6), 787–796.
- Chambers, R. A., Krystal, J. H., & Self, D. W. (2001). A neurobiological basis for substance abuse comorbidity in schizophrenia. *Society of Biological Psychiatry*, 50(2), 71–83.
- Chue P, Tibbo P, Wright E, Van Ens J. (2004). Client and community services satisfaction with an assertive community treatment subprogram for inner-city clients in Edmonton, Alberta. Canada *Journal of Psychiatry*, 49(9). <http://www1.cpaapc.org:8080/Publications/Archives/CJP/2004/september/chue.pdf>
- Conway, K. P., Compton, W., Stinson, F. S., & Grant, B. F. (2006). Lifetime comorbidity of DSM-IV mood and anxiety disorders and specific drug use disorders: Results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Journal of Clinical Psychiatry*, 67(2), 247–257.
- Derefinko KJ, Widiger TA. (2008). In Fatemi SH, Clayton PJ (Eds). *The Medical Basis of Psychiatry*, Third edition. Humana Press, New Jersey.
- Drake RE. (2007). Management of substance use disorder in schizophrenia patients: Current guidelines. *CNS Spectrums*, 12(10 Suppl 17), 27–32.

- Drake RE, Mueser KT, Brunette MF. (2007). Management of persons with co-occurring severe mental illness and substance use disorders: program implications. *World Psychiatry*, 6(3), 131–136.
- Drake RE, McHugo GJ, Xie H, Fox M, Packard J, Heimstetter B. (2006). Ten-year recovery outcomes for clients with co-occurring schizophrenia and substance use disorders. *Schizophrenia Bulletin*, 32(3), 464–473.
- Drake RE, Essock SM, Shaner A, Carey KB, Minkoff K, Kola L, Lynde D, Osher FC, Clark RE, Rickards L. (2001). Implementing dual diagnosis services for clients with severe mental illness. *Psychiatric Services*, 52(4), 469–476.
- Embry DD. (2002). The Good Behavior Game: a best practice candidate as a universal behavioral vaccine. *Clinical Children and Family Psychology Review*, 5(4), 273–297.
- First, M. B., Francis, A., & Pincus, H. A. (2002). *DSM-IV-TR Handbook of Differential Diagnosis*. Washington, DC: American Psychiatric Publishing, Inc.
- Gallagher, S. M., Penn, P. E., Brooks, A. J., & Feldman, J. (2006). Comparing the CAAPE, a new assessment tool for co-occurring disorders, with the SCID. *Psychiatric Rehabilitation Journal*, 30(1), 63–65.
- Goldstein, B. I., & Levitt, A. J. (2006). Factors associated with temporal priority in comorbid bipolar I disorder and alcohol use disorders: Results from the national epidemiologic survey on alcohol and related conditions. *Journal of Clinical Psychiatry*, 67(4), 643–649.
- Goldsmith, R. J., & Ries, R. K. (1988). Substance-induced mental disorders. In A. W. Graham, T. K. Schultz, & B. B. Wilford (Eds.), *Principles of Addiction Medicine* (2nd ed.) (pp. 969–981). Chevy Chase, MD: American Society of Addiction Medicine.
- Goldstein RB, Dawson DA, Saha TD, Ruan WJ, Compton WM, Grant BF. (2007). Antisocial behavioral syndrome and DSM-IV alcohol use disorders: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Alcohol Clinical Experimental Research*, 31(5), 814-828.
- Goodwin, R. D., Fergusson, D. M., & Horwood, L. J. (2004). Association between anxiety disorders and substance use disorders among young persons: Results of a 21-year longitudinal study. *Journal of Psychiatric Research*, 38(3), 295–304.
- Grant, B. F. (1996). Prevalence and correlates of drug use and DSM-IV drug dependence in the United States: Results of the National Longitudinal Alcohol Epidemiologic Survey. *Journal of Substance Abuse*, 8(2), 195–210.
- Grant, B. F., Stinson, F. S., Dawson, D. A., Chou, P., Dufour, M. C., Compton, W., et al. (2004). Prevalence and co-occurrence of substance use disorders and independent mood and anxiety disorders. *Archives of General Psychiatry*, 61, 807–816.
- Greenberg GA, Rosenheck RA. (2008). Jail incarceration, homelessness, and mental health: A national study. *Psychiatric Services*, 59(2), 170–177.
- Grella CE, Stein JA. (2006). Impact of program services on treatment outcomes of patients with comorbid mental and substance use disorders. *Psychiatric Services*, 57(7), 1007–1015.
- Hahn R, Fuqua-Whitney D, Wethington H et al.; CDC Task Force on Community Preventive Services. (2007). The effectiveness of universal schoolbased programs for the prevention of violent and aggressive behavior. *MMWR Recommended Report*, 56(RR-7), 1–12.
- Haning, W., Nunes, E., Penn, P.E. (2005). Response: The path to double recovery. *Science and Practice Perspectives*, 3(1), 22–24.
- Hasin, D., Samet, S., Nunes, E., Meydan, J., Matseoane, K., & Waxman, R. (2006). Diagnosis of comorbid psychiatric disorders in substance users assessed with the Psychiatric Research Interview for Substance and Mental Disorders for DSM-IV. *American Journal of Psychiatry*, 163(4), 689–696.

- Health Care for the Homeless Clinicians' Network, National Health Care for the Homeless Council. (2006). Integrating Primary & Behavioral Health Care for Homeless People. *Healing Hands*, 10(2), 1–6. <http://www.nhchc.org/Network/HealingHands/2006/May2006HealingHands.pdf>
- Herinckx H. (2008). Criminal Activity and Substance Abuse Study. Regional Research Institute for Human Services, Portland, OR. <http://www.nhchc.org/CriminalActivity&SubstAbuse.pdf>
- Hoffman, N. G. (2000). *CAAPE (Comprehensive Addiction and Psychological Evaluation) Manual*. Smithfield, RI: Evince Clinical Assessments.
- Johnson BA, Roache JD, Javors MA, DiClemente CC, Cloninger CR, Prihoda TJ, Bordnick PS, Ait-Daoud N, Hensler, J. (2000). Ondansetron for reduction of drinking among biologically predisposed alcoholic patients: a randomized controlled trial. *JAMA*, 284(8), 963–971.
- Kertesz SG, Mullins AN, Schumacher JE, Wallace D, Kirk K, Milby JB. (2007). Long-term housing and work outcomes among treated cocaine-dependent homeless persons. *Journal of Behavioral Health Service Research*, 34(1), 17–33.
- Kertesz SG, Madan A, Wallace D, Schumacher JE, Milby JB. (2006). Substance abuse treatment and psychiatric comorbidity: Do benefits spill over? Analysis of data from a prospective trial among cocaine-dependent homeless persons. *Substance Abuse Treatment, Prevention, and Policy*, 1(27), 1–8. <http://www.substanceabusepolicy.com/content/1/1/27>
- Kessler RC, Wang PS. (2008). The descriptive epidemiology of commonly occurring mental disorders in the United States. *Annual Review of Public Health*, 29, 115–129.
- Kessler, R. C., McGonagle, K. A., Zhao, S., Nelson, C. B., Hughes, M., Eshleman, et al. (1994). Lifetime and 12-month prevalence of DSM-II-R psychiatric disorders in the United States: Results from the National Comorbidity Survey. *Archives of General Psychiatry*, 51(1), 8–19.
- Komaromy M. (2009). The Addicted Brain. Presentation, Project ECHO Addiction Treatment Program, University of New Mexico.
- Komaromy M, O'Connell J. (2008). Treating Addiction with Medication. Presentation, National HCH Conference and Policy Symposium, Phoenix, AZ. <http://www.nhchc.org/2008conference/workshops/9.html>
- Kraybill K, Zerger S. (2003). *Providing Treatment for Homeless People with Substance Use Disorders: Case Studies of Six Programs*. National Health Care for the Homeless Council, Nashville, TN. <http://www.nhchc.org/Advocacy/FactSheets/CA05RCasestudies-FINAL5.pdf>
- Lamb HR, Weinberger LE, Marsh JS, Gross BH. (2007). Treatment prospects for persons with severe mental illness in an urban county jail. *Psychiatric Services*, 58(6), 782–786.
- Malcolm R, Myrick H, Roberts J, Wang W, Anton RF, Ballenger JC. (2002). The effects of carbamazepine and lorazepam on single versus multiple previous alcohol withdrawals in an outpatient randomized trial. *Journal of General Internal Medicine*, 17(5), 349–355.
- Mathalon, D. H., Pfefferbaum, A., Lim, K. O., Rosenbloom, M. J., & Sullivan, E. V. (2003). Compounded brain volume deficits in schizophrenia-alcoholism comorbidity. *Archives of General Psychiatry*, 60, 245–252.
- McGovern MP, Xie N, Segal SR, Siembab L, Drake RE. (2006). Addiction treatment services and co-occurring disorders: Prevalence estimates, treatment practices, and barriers. *Journal of Substance Abuse and Treatment*, 31(3), 267–275.
- Morse G. (1998). A Review of Case Management for People Who Are Homeless: Implications for Practice, Policy, and Research. Presented at the HUD/HHS Conference on Homelessness Research, October 1998: <http://aspe.hhs.gov/progsys/homeless/symposium/7-Casemgmt.htm>

- Mueser KT, Drake RE, Turner W, McGovern M. (2006). Comorbid substance use disorders and psychiatric disorders. In Miller WR, Carroll KM (Eds), *Rethinking Substance Abuse: What the Science Shows, and What We Should Do About It*. Guilford Press, New York.
- Mueser KT, Crocker AG, Frisman LB, Drake RE, Covell NH, Essock SM. (2006). Conduct disorder and antisocial personality disorder in persons with severe psychiatric and substance use disorders. *Schizophrenia Bulletin*, 32(4), 626–636.
- Mid-America Addiction Technology Transfer Center Network. (2008). Psychotherapeutic Medications 2008: What Every Counselor Should Know. ATTCN, Kansas City, MO, www.mattc.org
- Milby JB, Schumacher JE, Vuchinich RE, Freedman MJ, Kertesz S, Wallace D. (2008). Toward cost-effective initial care for substance-abusing homeless. *Journal of Substance Abuse Treatment*, 34, 180–191.
- Orson FM, Kinsey BM, Singh RA, Wu Y, Gardner T, Kosten TR. (2008). Substance abuse vaccines. *Annals of New York Academy of Science*, 1141, 257–269.
- Petras H, Kellam SG, Brown CH, Muthen BO, Ialongo NS, Poduska JM. (2008). Developmental epidemiological courses leading to antisocial personality disorder and violent and criminal behavior: Effects by young adulthood of a universal preventive intervention in first-and second-grade classrooms. *Drug and Alcohol Dependency*, 95 Suppl 1, S545–S559.
- Post R. (2009). 2007–2008 Alcohol & Drug Free Community Transitional Housing and Supported Employment Outcome Review. Central City Concern, Portland, OR.
- Quello, S. B., Brady, K. T., Sonne, S. C. (2005). Mood disorders and substance use disorder: A complex comorbidity. *Science and Practice Perspectives*, 3(1), 13–21.
- Regier, D. A., Narrow, W. E., Rae, D. S., Manderscheid, R. W., Locke, B. Z., & Goodwin, F. K. (1993). The de facto U.S. mental and addictive disorders service system: Epidemiologic Catchment Area prospective 1-year prevalence rates of disorders and services. *Archives of General Psychiatry*, 50, 85–94.
- Salekin RT. (2002). Psychopathy and therapeutic pessimism. Clinical lore or clinical reality? *Clinical Psychology Review*, 22(1) 79–112.
- Salloum IM, Cornelius JR, Daley DC, Kirisci L, Himmelhoch JM, Thase ME. (2005). Efficacy of valproate maintenance in patients with bipolar disorder and alcoholism: A double-blind placebo-controlled study. *Archives of General Psychiatry*, 62(1), 37–45.
- Samet, S., Nunes, E. V., & Hasin, D. (2004). Diagnosing comorbidity: Concepts, criteria, and methods. *Acta Neuropsychiatrica*, 16, 9–18.
- Shivani, R., Goldsmith, R. J., & Anthenelli, R. M. (2002). Alcoholism and psychiatric disorders: Diagnostic challenges. *Alcohol Research & Health*, 26, 90–98.
- Silver K. (2008). Co-occurring Disorders. Presentation, Group Work. Baltimore, MD.
- Substance Abuse and Mental Health Services Administration (SAMHSA), Center for Mental Health Services. (2008). Assertive Community Treatment Evidence-Based Practice Kit. <http://mentalhealth.samhsa.gov/cmhs/CommunitySupport/toolkits/community/>
- SAMHSA, Center for Substance Abuse Treatment. (2007). *Definitions and Terms Related to Co-Occurring Disorders*. COCE Overview Paper 1. DHHS Publication No. (SMA) 07–4163. Substance Abuse and Mental Health Services Administration (SAMHSA), Center for Mental Health Services, Rockville, MD. http://coce.samhsa.gov/cod_resources/PDF/OP1-DefinitionsandTerms-8-13-07.pdf
- SAMHSA, Office of Applied Studies. (2004). The NSDUH Report: *Adults with Co-Occurring Serious Mental Illness and a Substance Use Disorder*. US Department of Health and Human Services, Rockville, MD. <http://www.oas.samhsa.gov/2k4/coOccurring/coOccurring.htm>

- SAMHSA. (1999). Treatment Improvement Protocol (TIP) Series 34: Brief Interventions and Brief Therapies for Substance Abuse. <http://www.ncbi.nlm.nih.gov/books/bv.fcgi?rid=hstat5.chapter.59192>
- Tsemberis S, Eisenberg RF. (2000). Pathways to housing: Supported housing for street-dwelling homeless individuals with psychiatric disabilities. *Psychiatric Services*, 51(4), 487–493.
<http://psychservices.psychiatryonline.org/cgi/content/abstract/51/4/487>
- Widiger TA, Cadoret R, Hare R, Robins L, Rutherford M, Zanarini M, Alterman A, Apple M, Corbitt E, Forth A, Hart S, Kultermann J, Woody G, Frances A. (1996). DSM-IV antisocial personality disorder field trial. *Journal of Abnormal Psychology*, 105(1), 3–16.

These references were compiled from the following sources:

1. Health Care for the Homeless Clinicians' Network. Meeting the Challenges of Comorbid Mental Illness & Substance-Related Disorders. *Healing Hands* 13(1): 1–6.
<http://www.nhchc.org/Network/HealingHands/2009/Feb09HealingHands.pdf> [and other sources consulted in writing these articles]
2. SAMHSA SSA Co-Occurring Science Summary, 8-14-06.